

WE CLAIM:

1. An isolated polynucleotide encoding a receptor wherein said polynucleotide is selected  
5 from the group consisting of:

- a) a polynucleotide encoding a polypeptide having the deduced amino acid sequence  
of Figure 1 or a fragment, analog or derivative of said polypeptide; and
- b) a polynucleotide capable of hybridising to and which is at least 70% identical to  
10 the polynucleotide of Figure 1.

2. An isolated polynucleotide according to claim 1, wherein the polynucleotide is the  
polynucleotide of Figure 1.

3. An isolated polynucleotide according to claim 1, wherein the polynucleotide encodes a  
polypeptide having the deduced amino acid sequence of Figure 1 or a fragment, analog or  
derivative of said polypeptide.

4. An isolated polynucleotide comprising a region that encodes a variant of the  
polynucleotide of Figure 1, said variant sharing at least 95% amino acid identity with said  
Figure 1 polynucleotide

5. A recombinant DNA construct having incorporated therein a polynucleotide as defined in  
25 any one of claims 1 to 4.

6. A cell that has been selected to produce a receptor encoded by the polynucleotide as  
defined in any one of claims 1 to 4.

7. A cell according to claim 6 wherein said cell is genetically engineered to produce said receptor by incorporating expressibly therein a recombinant construct as defined in claim 5.

8. A cell according to claim 6 wherein said receptor is expressed endogenously.

9. A cell as defined in claim 6 which is a mammalian cell.

10. A receptor-binding membrane preparation derived from a cell as defined in claim 6.

11. A method of assaying a test ligand for binding with a receptor encoded by the polynucleotide as defined in any one of claims 1 to 4, which comprises the steps of incubating the test ligand under appropriate conditions with a receptor-producing cell as defined in claim 6, or with membrane preparation derived therefrom, and then determining whether binding between said receptor and said test ligand has occurred.

12. A method according to claim 11 wherein the binding between said receptor and said test ligand is determined by measuring a functional receptor response.

13. A method as defined in claim 12, wherein said functional receptor response is a second messenger response.

14. A method as defined in claim 13, wherein said second messenger is selected from the group consisting of intracellular cAMP and intracellular calcium ion.

15. A receptor encoded by the polynucleotide as defined in any one of claims 1 to 4, in an isolated form essentially free from other proteins of human origin.

16. A ligand-binding fragment of a receptor encoded by the polynucleotide defined in any one of claims 1 to 4.

17. An antibody which binds a mammalian receptor encoded by the polynucleotide defined in any one of claims 1 to 4.

5 18. An immunogenic fragment of a human receptor wherein said receptor is encoded by the polynucleotide defined in any one of claims 1 to 4.

19. An oligonucleotide which comprises at least about 17 nucleic acids and which selectively hybridizes with a polynucleotide defined in claim 1 or complement thereof.

10  
20  
30  
40  
50  
60  
70  
80  
90  
100  
110  
120  
130  
140  
150  
160  
170  
180  
190  
200  
210  
220  
230  
240  
250  
260  
270  
280  
290  
300  
310  
320  
330  
340  
350  
360  
370  
380  
390  
400  
410  
420  
430  
440  
450  
460  
470  
480  
490  
500  
510  
520  
530  
540  
550  
560  
570  
580  
590  
600  
610  
620  
630  
640  
650  
660  
670  
680  
690  
700  
710  
720  
730  
740  
750  
760  
770  
780  
790  
800  
810  
820  
830  
840  
850  
860  
870  
880  
890  
900  
910  
920  
930  
940  
950  
960  
970  
980  
990  
1000